

ENVIRONMENT

Biodiversity: “Today’s low rates of marine extinction may be the prelude to a major extinction pulse” in oceans

Sunday 18 January 2015, by [Science](#), [ZIMMER Carl](#) (Date first published: 15 January 2015).

Contents

- [Marine defaunation: Animal \(...\)](#)
- [Ocean Life Faces Mass Extincti](#)

Marine defaunation: Animal loss in the global ocean

REVIEW

Douglas J. McCauley*, Malin L. Pinsky, Stephen R. Palumbi, James A. Estes, Francis H. Joyce, Robert R. Warner

For Author Affiliations see the original

SUMMARY

BACKGROUND

Comparing patterns of terrestrial and marine defaunation helps to place human impacts on marine fauna in context and to navigate toward recovery. Defaunation began in earnest tens of thousands of years later in the oceans than it did on land. Although defaunation has been less severe in the oceans than on land, our effects on marine animals are increasing in pace and impact. Humans have caused few complete extinctions in the sea, but we are responsible for many ecological, commercial, and local extinctions. Despite our late start, humans have already powerfully changed virtually all major marine ecosystems.

ADVANCES

Humans have profoundly decreased the abundance of both large (e.g., whales) and small (e.g., anchovies) marine fauna. Such declines can generate waves of ecological change that travel both up and down marine food webs and can alter ocean ecosystem functioning. Human harvesters have also been a major force of evolutionary change in the oceans and have reshaped the genetic structure of marine animal populations. Climate change threatens to accelerate marine defaunation over the next century. The high mobility of many marine animals offers some increased, though limited, capacity for marine species to respond to climate stress, but it also exposes many species to increased risk from other stressors. Because humans are intensely reliant on ocean ecosystems for food and other ecosystem services, we are deeply affected by all of these forecasted changes.

Three lessons emerge when comparing the marine and terrestrial defaunation experiences: (i) today's low rates of marine extinction may be the prelude to a major extinction pulse, similar to that observed on land during the industrial revolution, as the footprint of human ocean use widens; (ii) effectively slowing ocean defaunation requires both protected areas and careful management of the intervening ocean matrix; and (iii) the terrestrial experience and current trends in ocean use suggest that habitat destruction is likely to become an increasingly dominant threat to ocean wildlife over the next 150 years.

OUTLOOK

Wildlife populations in the oceans have been badly damaged by human activity. Nevertheless, marine fauna generally are in better condition than terrestrial fauna: Fewer marine animal extinctions have occurred; many geographic ranges have shrunk less; and numerous ocean ecosystems remain more wild than terrestrial ecosystems. Consequently, meaningful rehabilitation of affected marine animal populations remains within the reach of managers. Human dependency on marine wildlife and the linked fate of marine and terrestrial fauna necessitate that we act quickly to slow the advance of marine defaunation.

Timeline (log scale) of marine and terrestrial defaunation.

The marine defaunation experience is much less advanced, even though humans have been harvesting ocean wildlife for thousands of years. The recent industrialization of this harvest, however, initiated an era of intense marine wildlife declines. If left unmanaged, we predict that marine habitat alteration, along with climate change (colored bar: IPCC warming), will exacerbate marine defaunation.

* *Science* 16 January 2015; Vol. 347 no. 6219 DOI: 10.1126/science.1255641:
<http://www.sciencemag.org/content/347/6219/1255641>

Ocean Life Faces Mass Extinction, Broad Study Says

A team of scientists, in a groundbreaking analysis of data from hundreds of sources, has concluded that humans are on the verge of causing unprecedented damage to the oceans and the animals living in them.

"We may be sitting on a precipice of a major extinction event," said Douglas J. McCauley, an ecologist at the University of California, Santa Barbara, and an author of the new research, which was published on Thursday in the journal *Science* (see above).

But there is still time to avert catastrophe, Dr. McCauley and his colleagues also found. Compared with the continents, the oceans are mostly intact, still wild enough to bounce back to ecological health.

"We're lucky in many ways," said Malin L. Pinsky, a marine biologist at Rutgers University and another author of the new report. "The impacts are accelerating, but they're not so bad we can't reverse them."

Scientific assessments of the oceans' health are dogged by uncertainty: It's much harder for researchers to judge the well-being of a species living underwater, over thousands of miles, than to track the health of a species on land. And changes that scientists observe in particular ocean ecosystems may not reflect trends across the planet.

Dr. Pinsky, Dr. McCauley and their colleagues sought a clearer picture of the oceans' health by pulling together data from an enormous range of sources, from discoveries in the fossil record to statistics on modern container shipping, fish catches and seabed mining. While many of the findings already existed, they had never been juxtaposed in such a way.

A number of experts said the result was a remarkable synthesis, along with a nuanced and encouraging prognosis.

"I see this as a call for action to close the gap between conservation on land and in the sea," said Loren McClenachan of Colby College, who was not involved in the study.

There are clear signs already that humans are harming the oceans to a remarkable degree, the scientists found. Some ocean species are certainly overharvested, but even greater damage results from large-scale habitat loss, which is likely to accelerate as technology advances the human footprint, the scientists reported.

Coral reefs, for example, have declined by 40 percent worldwide, partly as a result of climate-change-driven warming.

Some fish are migrating to cooler waters already. Black sea bass, once most common off the coast of Virginia, have moved up to New Jersey. Less fortunate species may not be able to find new ranges. At the same time, carbon emissions are altering the chemistry of seawater, making it more acidic.

"If you cranked up the aquarium heater and dumped some acid in the water, your fish would not be very happy," Dr. Pinsky said. "In effect, that's what we're doing to the oceans."

Fragile ecosystems like mangroves are being replaced by fish farms, which are projected to provide most of the fish we consume within 20 years. Bottom trawlers scraping large nets across the sea floor have already affected 20 million square miles of ocean, turning parts of the continental shelf to rubble. Whales may no longer be widely hunted, the analysis noted, but they are now colliding more often as the number of container ships rises.

Mining operations, too, are poised to transform the ocean. Contracts for seabed mining now cover 460,000 square miles underwater, the researchers found, up from zero in 2000. Seabed mining has the potential to tear up unique ecosystems and introduce pollution into the deep sea.

The oceans are so vast that their ecosystems may seem impervious to change. But Dr. McClenachan warned that the fossil record shows that global disasters have wrecked the seas before. "Marine species are not immune to extinction on a large scale," she said.

Until now, the seas largely have been spared the carnage visited on terrestrial species, the new analysis also found.

The fossil record indicates that a number of large animal species became extinct as humans arrived on continents and islands. For example, the moa, a giant bird that once lived on New Zealand, was wiped out by arriving Polynesians in the 1300s, probably within a century.

But it was only after 1800, with the Industrial Revolution, that extinctions on land really accelerated.

Humans began to alter the habitat that wildlife depended on, wiping out forests for timber, plowing under prairie for farmland, and laying down roads and railroads across continents.

Species began going extinct at a much faster pace. Over the past five centuries, researchers have recorded 514 animal extinctions on land. But the authors of the new study found that documented extinctions are far rarer in the ocean.

Before 1500, a few species of seabirds are known to have vanished. Since then, scientists have documented only 15 ocean extinctions, including animals such as the Caribbean monk seal and the Steller's sea cow.

While these figures are likely underestimates, Dr. McCauley said that the difference was nonetheless revealing.

"Fundamentally, we're a terrestrial predator," he said. "It's hard for an ape to drive something in the ocean extinct."

Many marine species that have become extinct or are endangered depend on land — seabirds that nest on cliffs, for example, or sea turtles that lay eggs on beaches.

Still, there is time for humans to halt the damage, Dr. McCauley said, with effective programs limiting the exploitation of the oceans. The tiger may not be salvageable in the wild — but the tiger shark may well be, he said.

"There are a lot of tools we can use," he said. "We better pick them up and use them seriously."

Dr. McCauley and his colleagues argue that limiting the industrialization of the oceans to some regions could allow threatened species to recover in other ones. "I fervently believe that our best partner in saving the ocean is the ocean itself," said Stephen R. Palumbi of Stanford University, an author of the new study.

The scientists also argued that these reserves had to be designed with climate change in mind, so that species escaping high temperatures or low pH would be able to find refuge.

"It's creating a hopscotch pattern up and down the coasts to help these species adapt," Dr. Pinsky said.

Ultimately, Dr. Palumbi warned, slowing extinctions in the oceans will mean cutting back on carbon emissions, not just adapting to them.

"If by the end of the century we're not off the business-as-usual curve we are now, I honestly feel there's not much hope for normal ecosystems in the ocean," he said. "But in the meantime, we do have a chance to do what we can. We have a couple decades more than we thought we had, so let's please not waste it."

Carl Zimmer

*

http://www.nytimes.com/2015/01/16/science/earth/study-raises-alarm-for-health-of-ocean-life.html?_r=1

A version of this article appears in print on January 16, 2015, on page A1 of the New York edition with the headline: Ocean Life Faces Mass Extinction, Broad Study Says.

